


ENGRAFTABLE NEURAL PROGENITOR & STEM CELLS FOR BRAIN TUMOR THERAPY

FIELD OF THE INVENTION

5

This invention is in the field of gene therapy, more particularly the field of using neuronal cells to treat brain tumors.

10



This application is a continuation-in-part of pending U.S. Serial No. 09/133,873, ^{now U.S. Patent No. 5,958,767} filed on August 14, 1998, which is incorporated herein by reference.

Government Support

15

This invention was made with support from the NIH under grant number P20-HD18655, and the United States government has certain rights in this invention.

20

BACKGROUND

An effective gene therapy for the treatment of brain tumors has been an elusive goal for many years. Glioblastoma multiforma, which is virtually untreatable, and the less malignant anaplastic astrocytoma account for about one-quarter of the 5,000 intracranial gliomas diagnosed yearly in the United States; 75 percent of gliomas in adults are of this category. Because of its profound and uniform morbidity, it contributes more to the cost of cancer on a per capita basis than does any other tumor. The patient, commonly stricken in the fifth decade of life, enters a cycle of repetitive hospitalizations and operations while experiencing the progressive complications associated with relatively ineffective treatments of radiation and chemotherapy ("Harrison's Principles of Internal

Medicine,” edited by Isselbacher, Braunwald, Wilson, Martin, Fauci and Kasper, 13th Edition, p.2262, McGraw-Hill, Inc. 1994).

One of the impediments to gene therapy of brain tumors such as gliomas, has been the degree to which they expand, migrate widely and infiltrate normal tissue. Most gene therapy strategies to date are viral vector-based, yet extensive distributions of sufficient amounts of viral vector-mediated genes to large regions and numbers of cells typically in need has often been disappointingly limited. Interestingly, one of the defining features of normal neural progenitors and stem cells is their migratory quality. Neural stem cells (NSCs) are immature, uncommitted cells that exist in the developing, and even adult, CNS and postulated to give rise to the array of more specialized cells of the CNS. They are operationally defined by their ability to self-renew and to differentiate into cells of most (if not all) neuronal and glial lineages in multiple anatomical & development contexts, and to populate developing and /or degenerating CNS regions.¹⁻⁵

With the first recognition that neural cells with stem cell properties, reproduced in culture, could be reimplanted into mammalian brain where they could reintegrate appropriately and seamlessly in the neural architecture and stably express foreign genes⁶⁻⁷, gene therapists began to speculate how such a phenomenon might be harnessed for therapeutic purposes. These, and the studies which they spawned (reviewed elsewhere^{1-5,8}), provided hope that the use of neural progenitor/stem cells, by virtue of their inherent biology, might circumvent some of the present limitations of presently available gene transfer vehicles (e.g., non-neural cells, viral vectors, synthetic pumps), and provide the basis for a variety of novel therapeutic strategies.

Their use as graft material has been clearly illustrated by the prototypical neural progenitor clone, C17.2, a clone with which we have had extensive experience^{6,9-16,17} and which was used in the studies presented here. C17.2 is a mouse cell line from postnatal day 0 cerebellum immortalized by infection with a retroviral construct containing the avian myc gene. This line has been transduced to constitutively express the *lacZ* and *neoR* genes. When transplanted into germinal zones throughout the brain, these cells have been shown to migrate, cease dividing, and participate in the normal development of multiple regions at multiple stages (fetus to adult) along the murine neuraxis,

differentiating appropriately into diverse neuronal and glial cell types as normal, non-tumorigenic cytoarchitectural constituents. They intermingle non-disruptively with endogenous neural progenitor/stem cells, responding to the same spatial and temporal cues in a similar manner. Crucial for therapeutic considerations, the structures to which C17.2 cells contribute develop and maintain neuroanatomical normality. In their earliest therapeutic use, they served to deliver a missing gene product throughout the brains of mice with a lysosomal deficiency state and cross-corrected host cells by release and uptake of a lysosomal enzyme⁹. The feasibility of a neural progenitor/stem cell-based strategy for the delivery of therapeutic molecules directly to and throughout the CNS was first affirmed by correcting the widespread neuropathology of a murine model of the genetic neurodegenerative lysosomal storage disease mucopolysaccharidosis type VII, caused by an inherited deletion of the β -glucuronidase (GUSB) gene, a condition that causes mental retardation and early death in humans. Exploiting their ability to engraft diffusely and become integral members of structures throughout the host CNS, GUSB-secreting NSCs were introduced at birth into subventricular germinal zone, and provided correction of lysosomal storage in neurons and glia throughout mutant brains. In so doing, it established that neural transplantation of neural progenitor cells could provide a novel therapeutic modality.

What is needed is a way to treat tumors which are diffuse, infiltrating and/or metastasizing. What is needed is a way to treat tumors locally to maximize the impact on the tumor and reduce the toxicity to the patient.

SUMMARY OF THE INVENTION

An isolated pluripotent neuronal cell having the capacity to differentiate into at least different types of nerve cells is disclosed. The pluripotent cell is further characterized by having a migratory capacity whereby the cell is capable of travelling from a first location where the neuronal cell is administered to a second location at which there is at least one tumor cell, having the ability to travel through and around a tumor, whereby a plurality of the neuronal cells are capable of surrounding the tumor; and

having the capacity to track at least one infiltrating tumor cell, thereby treating infiltrating and metastasizing tumors.

The neuronal cell may be an isolated neural stem cell. The neuronal cell is optionally treated to secrete a cytotoxic substance. The neuronal cell alternatively is transformed with factors that directly promote differentiation of neoplastic cells. Alternatively, the neuronal cell is transformed with viral vectors encoding therapeutic genes to be incorporated by tumor cells. In another embodiment, the neuronal cell can be transformed with viral vectors encoding suicide genes, differentiating agents, or receptors to trophins to be incorporated into tumor cells. The neuronal cells if administered on the same side or a contralateral side of the brain from the tumor, are capable of reaching the tumor.

In another embodiment there is provided a method of converting a migrating neuronal cell to a migrating packaging/producer cell, said method includes the steps of a) providing a neuronal cell which constitutively produces a marker such as β -gal; b) cotransfecting the neuronal cell with an amphotropic pPAM3 packaging plasmid and a puromycin selection plasmid pPGKpuro; c) selecting transfected cells in puromycin; d) selecting for cell surface expression of the amphotropic envelope glycoprotein coat; e) isolating cells by fluorescent activated cell sorting using monoclonal antibody 83A25; and f) screening the cells of step e for their packaging ability by assessing which colonies packaged lacZ into infectious viral particles. Thus there is produced a migratory neuronal cell capable of being transfected with a gene of choice, so that viral particles expressing the gene of choice are produced and disseminated over a wide area of the central nervous system by a plurality of the transfected packaging cells.

The method of converting the migratory neuronal cell into a packaging cell line wherein step f is performed by a virus focus assay for β -gal production. Alternatively the method can be performed with a prodrug activation enzyme as the gene of choice. Alternatively, the prodrug activation enzyme is *E.coli* cytosine deaminase (CD), HSV-TK or cytochrome p450. More preferably, the prodrug activation enzyme is *E.coli* cytosine deaminase (CD).

Also disclosed is a novel cell packaging line for the central nervous system. The cell line includes neuronal cells which constitutively produce a marker such as β -gal, have been cotransfected with an amphotropic pPAM3 packaging plasmid and a puromycin selection plasmid pPGKpuro; are selected in puromycin, for cell surface expression of the amphotropic envelope glycoprotein coat and for fluorescence using monoclonal antibody 83A25, and for their packaging ability by assessing which colonies packaged lacZ into infectious viral particles. The resulting cells are capable of packaging and releasing particles or vectors which, in turn, may serve as vectors for gene transfer to central nervous system cells. The particles in the novel cell packaging line can be replication-defective retroviral particles. The vectors in the novel cell packaging line can be replication-conditional herpes virus vectors.

BRIEF DESCRIPTION OF DRAWINGS

Figures 1A and 1B illustrate the migratory capacity of neural progenitor/stem C17.2 cells *in vitro*. After 5 days of incubation there was a wide distribution of C17.2 cells (Fig. 1B), suggesting that they had migrated far from their initial seeding in the cylinder, compared to TR-10 cells (Fig. 1A), which remained localized to the area of initial seeding in the cylinders. These patterns were observed whether the cells were plated directly on top of the glioma cells (right-sided cylinder [arrows]) or simply in juxtaposition to them (center cylinder [arrows]).

Figures 2A, 2B, 2C and 2D illustrate foreign gene-expressing neural progenitor/stem cells extensive migration throughout experimental tumor mass, and slightly beyond advancing tumor edge, appearing to “track” migrating tumor cells. (Fig. 2A) day 2 shown at 4X; arrowheads demarcate the approximate edges of tumor mass; (Fig. 2B) high power at 10X where Xgal, blue-staining NSCs [arrows] are interspersed between tumor cells staining dark red. (Fig. 2C) View of tumor mass 10 days after intratumoral injection showing Xgal+blue, C17.2 NSCs have infiltrated the tumor but largely stop at the edge of the darkly red stained tumor tissue with some migration into surrounding tissue when the blue-staining NSC appears to be “following” an invading, “escaping” cell [arrow] (10X). (Fig. 2D) CNS-1 tumor cells implanted into an adult

nude mouse frontal cortex, there is extensive migration and distribution of blue C17.2 cells throughout the infiltrating experimental tumor bed, up to and along the infiltrating tumor edge [arrows], and where many tumor cells are invading normal tissue, into surrounding tissue in virtual juxtaposition to aggressive tumor cells [arrows] (10X).

Figures 3A, 3B, 3C, 3D, 3E, 3F, 3G and 3H illustrate the neural progenitor/stem cells appearance to "track" migrating tumor cells away from main tumor mass; (Figs. 3A, 3B) parallel sections: low power C17.2 cells distributed throughout tumor and surrounding edge [Fig. 3A) Xgal and neutral red, Fig. 3B) double immunofluorescent labelling with texas red and FITC]; (Figs. 3C, 3D) low and high power of tumor edge and migrating tumor cell in juxtaposition to C17.2 cell (Xgal and neutral red); (Figs. 3G, 3H) low and high power of single migrating tumor cells in juxtaposition to C17.2 cells (double immunofluorescent labelling with texas red and FITC).

Figures 4A, 4B, 4C, 4D, 4E, 4F and 4G illustrate neural progenitor/stem cells implanted at distant site from main tumor bed migrating throughout normal tissue target CNS-1 tumor cells; (Figs. 4A, 4B) same hemisphere: 3×10^4 CNS-1 tumor cells implanted into right frontal lobe. On day 6, 4×10^4 C17-2 cells injected into right frontoparietal lobe (4mm caudal tumor injection). Animals sacrificed on day 12 (shown) and day 21, C17-2 cells seen in tumor bed (Xgal and neutral red). (Figs. 4C, 4D, 4E) Contralateral hemisphere: 3×10^4 CNS-1 tumor cells implanted into left frontal lobe and 5×10^4 CNS-1 tumor cells implanted into left frontoparietal lobe. On day 6, 8×10^4 C17-2 cells were injected into right front lobe. Animals were sacrificed on day 12 and 21 (shown); c) 4x C17.2 cells (red) seen migrating towards tumor (green) from opposite side of the brain, d) 10x C17.2 cells (red) seen actively migrating across central commissure (double immunofluorescence), e) 20x C17-2 cells (blue) seen entering tumor (black arrows) (Xgal/ neutral red). (Figs. 4F, 4G) Intraventricular: 5×10^4 CNS-1 tumor cells were implanted into right frontal lobe. On day 6, 8×10^4 C17.2 cells were injected into right or left (shown) lateral ventricle.

DETAILED DESCRIPTION OF INVENTION

The experiments presented herein demonstrate that NSCs (prototypical clone C17.2) when implanted into an experimental glioma, will distribute throughout the tumor and migrate along with aggressively advancing tumor cells, while continuing to express their reporter gene lacZ. (One of the glioma lines used, astrocytoma cell line CNS-1, demonstrates single cell infiltration and invasive characteristics similar to those of human glioblastomas¹⁸). Furthermore, the neural progenitor/stem cells seem to migrate slightly beyond and surround the invading tumor border. In additional experiments, where neural progenitors were implanted at a distant site from the tumor bed, in the same hemisphere, opposite hemisphere, or lateral ventricle, they migrated through normal tissue moving specifically toward CNS-1 tumor cells. They were found to accumulate in or near the tumor bed as well as near or in direct juxtaposition to the individual infiltrating tumor cells.

Not wishing to be bound by any particular theory, the inventors propose that this neural progenitor/stem cell system migrate towards a trophic gradient of growth factors produced by the tumor cells. Thus, NSCs may provide a unique platform for the dissemination of therapeutic genes to the proximity of or into tumors that previously were inaccessible. These observations further suggest a number of other new gene therapy approaches. These may include the dissemination of cytotoxic gene products, but could also include factors that directly promote differentiation of neoplastic cells as well as the more efficacious delivery of viral vectors encoding therapeutic genes to be incorporated by tumor cells (e.g. suicide genes, differentiating agents, receptors to trophins). Because NSCs can be engineered to package and release replication-defective retroviral particles or replication-conditional herpes virus vectors which, in turn, may serve as vectors for the transfer of genes to CNS cells, neural progenitor/stem cells should serve to magnify the efficacy of viral-mediated gene delivery to large regions in the brain.

One effective mode of therapy for experimental brain tumors has been prodrug activation. Initially, prodrug activation enzymes were limited to antibodies directed against tumor enriched antigens. New strategies incorporate genes for these enzymes into viral vectors. Among the prodrug activating systems shown to be effective for gliomas

E. coli cytosine deaminase (CD), HSV-TK and cytochrome p450 have been demonstrated to have a drug mediated bystander effect. Of these CD gives the best reported “bystander” effect. CD converts the nontoxic prodrug 5-fluorocytosine (5-FC) to 5-fluorouridine (5-FU) metabolites. 5-FU is a chemotherapeutic agent which has selective toxicity for actively dividing cells, thus primarily targeting tumor cells. In addition, 5-FU and its toxic metabolites can readily pass into adjacent and surrounding cells by non-facilitated diffusion. Brain tumors may require only a small number of cells expressing CD (about 2% evenly distributed) to generate significant anti-tumor effects when treated with systemic, non-toxic levels of 5-FC. Our results support the hypothesis that transduced NSCs would disperse CD expression efficiently throughout the tumor and even “track” single migrating, “escaping” tumor cells.

Another approach to brain tumor gene therapy has been selective gene transfer to tumor cells in combination with pharmacotherapy, e.g., the HSV-TK gene, when transduced via retrovirus into a dividing population of brain tumor cells, confers a lethal sensitivity to the drug ganciclovir. Recent modifications of retroviral constructs to increase efficiency of infection and cell-specific targeting hold promise for enhancing the potency of this strategy. Again, through the “bystander effect”, tumor destruction is effective even when only a fraction of the cells express HSV-TK; adjacent tumor cells not expressing HSV-TK also appear to be eliminated. Attempts to improve efficiency of tumor destruction have focused on increasing the number of cells expressing the HSV-TK gene. The use of NSCs as packaging cells (which might then be self-eliminated) may prove to be an effective extended delivery system of the lethal gene to neighboring mitotic tumor cells, especially individual, infiltrating tumor cells.

In conclusion, genetically modified neural progenitor/stem cells have the potential to supply a range of tumor selective agents throughout mature and developing brains. The experiments presented here demonstrate the ability of NSCs: (1) to migrate/distribute quickly and effectively throughout the main tumor bed when implanted directly into the experimental gliomas; (2) to migrate slightly beyond and “surround” (as if to contain) the invading tumor border; (3) to seemingly “track” individual, infiltrating tumor cells into surrounding tissue; (4) to migrate through normal tissue from distant sites to target CNS-

1 tumors; and (5) to show stable expression of a foreign gene, in this case *lacZ*,
 throughout the tumor bed and in juxtaposition to tumor cells. These results lay the
 groundwork for future therapeutic brain tumor studies, providing critical support for the
 use of neural progenitor/stem cells as an effective delivery vehicle for tumor directed,
 5 vector-mediated enzyme/prodrug gene therapy.

Other cells

The HCN-1 cell line is derived from parental cell lines from the cortical tissue of
 a patients with unilateral megalencephaly growth (Ronnett G.V. et al. Science 248:603-5,
 1990). HCN-1A cells have been induced to differentiate to a neuronal-like morphology
 10 and stain positively for neurofilament, neuron-specific enolase and p75NGFR, but not for
 myelin basic protein, S-100 or glial fibrillary acidic protein (GFAP). Because these cells
 also stain positively for γ -amino butyric acid and glutamate, they appear to become
 neuro-transmitting bodies. Earlier Poltorak M et al. (Cell Transplant 1(1):3-15, 1992)
 observed that HCN-1 cells survived in the brain parenchyma and proposed that these cells
 15 may be suitable for intracerebral transplantation in humans.

Ronnet GV et al. (Neuroscience 63(4):1081-99, 1994) reported that HCN-1 cells
 grew processes resembling neurons when exposed to nerve growth factor, dibutyl
 cyclic AMP and isobutylmethylxanthine.

The nerve cells also can be administered with macrophages which have been
 20 activated by exposure to peripheral nerve cells. Such activated macrophages have been
 shown to clean up the site of CNS trauma, for example a severed optic nerve, after which
 new nerve extensions started to grow across the lesion. Implanting macrophages exposed
 to CNS tissue (which secretes a chemical to inhibit macrophages) or nothing at all
 resulted in little or no regeneration (Lazarov-Spiegler et al. FASEB J. 10:1, 1996).

25 Fetal pig cells have been implanted into patients with neurodegenerative diseases,
 such as Parkinson's disease and Huntington's chorea, and intractable seizures, in whom
 surgical removal of the excited area would otherwise have been performed. Such cells, if
 properly screened for retroviruses, could also be used in the inventive method.

Neural crest cells are isolated and cultured according to Stemple and Anderson
 30 (U.S. Patent No. 5,654,183), which is incorporated herein by reference, with the

modification that basic fibroblast growth factor (bFGF) is added to the medium at concentrations ranging from 5 to 100 ng/ml in 5 ng/ml increments. Neural crest cells so cultured are found to be stimulated by the presence of FGF in increasing concentrations about 1 or 5 ng/ml. Such cells differentiate into peripheral nerve cells, which can be used in the instant invention.

Other cytokines, growth factors and drugs

Certain cytokines, growth factors and drugs are optionally used in the transplant area or may be administered concomitantly with the transplant.

Known cytokines include interleukins (IL) IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-10, and IL-11; tissue necrosis factors (TNF), TNF α , also lymphotoxin (LT) and TNF β ; interferons (IFN) IFN α , IFN β and IFN γ ; and tissue growth factor (TGF). The colony-stimulating factors (CSFs) are specific glycoproteins that are thought to be involved in the production, differentiation and function of stem cells.

Nerve growth factor (NGF) has been shown to increase the rate of recovery in spatial alternation tasks after entorhinal lesions, possibly by acting on cholinergic pathways (Stein and Will, Brain Res. 261:127-31, 1983).

EXAMPLES

Experimental Methods

Cells:

C17-2 and TR-10 cells were maintained in Dulbecco's modified Eagle's medium (DMEM; Mediatech, Washington, DC) supplemented with 10% fetal calf serum (FCS; Sigma, St. Louise, MO), 5% horse serum (HS; Gibco), 1% Glutamine (2mM; Gibco), 1% penicillin/streptomycin (Sigma). CNS-1 cells were stably transduced with the PGK-GFP-IRES-NeoR retroviral vector construct to express green fluorescent protein (GFP) as previously described (ref. Aboody-Guterman et. al, 1997), and maintained in RPMI-1640 (Bio Whittaker) supplemented with 10% FCS and 1% penicillin/streptomycin (Sigma). Cell structure studies were performed in 100mm petri dishes under standard conditions: humidified, 37°C, 5% CO₂ incubator. *In vitro* studies; CNS-1 glioma cells were plated to

approximately 60-70% confluency around a 5mm cylinder (i.e. free of CNS-1 cells) into which 40,000 C17.2 or TR-10 cells plated overnight. At the same time, 40,000 C17.2 or TR-10 cells were placed into a 5mm cylinder placed directly on top of adhered CNS-1 cells. The next day, cylinders were removed and plates rinsed well with PBS to remove any floating cells, media was replaced, and plates incubated for 5 days. Plates were subsequently stained for β -galactosidase overnight after .5% glutaraldehyde fixation. (Note: both C17.2 and TR-10 cells are >90% blue with X-gal staining). In vivo studies; 48 hours prior to transplant, C17.2 and TR-10 cells were incubated with BUdR (Sigma) at a concentration of 10 μ M. Plated cells were rinsed with PBS, trypsinized, resuspended in media and counted on the Coulter counter. Desired number of cells were spun down at 4°C in the centrifuge for 4 minutes and 1100 rpm to obtain a pellet. Media was removed; cells were rinsed by resuspending in PBS and respun. PBS was removed and the appropriate amount of PBS added to resuspend cells at final desired concentration. Cells were kept on ice, and gently triturated prior to each animal injection. Cells not labelled with BUdR were prepared for injection in similar manner.

Animals:

Animal studies were performed in accordance with guidelines issued by the Massachusetts General Hospital Subcommittee on Animal Care. Animals used: adult CD-Fisher rats (Charles River) and 8-10 week old adult, approximately 20 gram female nude mice (random bred Swiss white obtained from Cox 7, MGH-East).

Surgery and Sacrifice:

Animals were anesthetized by an i.p. injection of .15 ml of 20% ketamine HCL (KETALAR 100 mg/ml; Parke-Davis, Morris Plains, NJ), 20% xylazine (ROMPUN 20 mg/ml; Miles Inc., Shawnee Mission, KS), 60% sodium chloride (0.9%; Abbott Laboratories, North Chicago, IL) and immobilized in stereotactic apparatus (Kopf, Tujunga, CA). Intracerebral injections were stereotactically performed by making a linear scalpel skin incision on top of the skull. A burr hole was drilled into the skull with a high speed drill 2mm lateral to the bregma on the coronal suture. After incising the dura with a sterile needle and obtaining hemostasis, desired number of tumor cells suspended in 1 μ l of 1X Dulbecco's phosphate-buffered salt solution (PBS pH 7.4;

Mediatech, Herndon, VA) were injected with a 26 gauge 5 μ l Hamilton syringe to specified location (see protocols below) over a 3 to 5 minute period. After retracting the needle over a 2-4 minute period, bone-wax (Ethicon, Somerville, NJ) was used to occlude the burr hole, betadine applied to surgical area, and the skin sutured closed. Animals receiving a second injection at a later date were anesthetized, immobilized in stereotactic apparatus, and cells injected as per specific protocol (see below). Animals were sacrificed on stated days with an overdose of anesthesia and subsequent intracardiac perfusion with PBS followed by 4% paraformaldehyde+2mM $MgCl_2$ (pH 7.4). Brains were removed and post-fixed overnight at 4 $^{\circ}C$ and then transferred to 30% sucrose in PBS + 2mM $MgCl_2$ (pH 7.4) for 3-7 days to cryoprotect. Brains were stored at -80 $^{\circ}C$ and then 10-15 micron coronal serial sections were cut to cryostat (Leica CM 3000).

BUDR labelling of engrafted C17-2 cells:

Selected animals received 3 intraperitoneal injections of 1 ml/100g body weight 20uM BUDR stock solution (Sigma) over 24 hours prior to sacrifice (.2 ml/injection per 20g mouse).

Histopathological and Immunohistochemical Studies:

Tissue sections were stained with (1) X-gal and counterstained with neutral red (2) hematoxylin and eosin (3), double immunofluorescent labelling was performed with texas red anti-beta-galactosidase and FITC anti-GFP. Slides were examined with light microscopy, fluorescent microscopy. CNS-1 tumor cells were also examined without staining under confocal fluorescent microscopy.

Example 1. Migratory Capacity of NSCs in Culture

To determine properties of the NSCs in association with glioma cells, studies were initially performed in culture comparing the relative migratory capacity of NSCs (clone C17.2) to fibroblasts (the lacZ-expressing TR-10 fibroblast cell line) when co-cultured with glioma cells. C17.2 and TR-10 cells were maintained in Dulbecco's modified Eagle's medium (DMEM; Mediatech, Washington, DC) supplemented with 10% fetal calf serum (FCS; Sigma, St. Louise, MO), 5% horse serum (HS; Gibco), 1% Glutamine (2mM; Gibco), 1% penicillin/streptomycin (Sigma). CNS-1 cells were stably

transduced with the PGK-GFP-IRES-NeoR retroviral vector construct to express green fluorescent protein (GFP) as previously described (ref. Aboody-Guterman et. al, 1997), and maintained in RPMI-1640 (Bio Whittaker) supplemented with 10% FCS and 1% penicillin/streptomycin (Sigma). Cell structure studies were performed in 100mm petri dishes under standard conditions: humidified, 37 ° C, 5% CO2 incubator. CNS-1 glioma cells were plated to approx. 60-70% confluency around a 5mm cylinder (i.e. free of CNS-1 cells) into which 40,000 C17.2 or TR-10 cells plated overnight. At the same time, 40,000 C17.2 or TR-10 cells were placed into a 5mm cylinder placed directly on top of adhered CNS-1 cells. The next day, cylinders were removed and plates rinsed well with

10 PBS to remove any floating cells, media was replaced, and plates incubated for 5 days. Plates were subsequently stained for β -galactosidase overnight after .5% glutaraldehyde fixation. (Note: both C17.2 and TR-10 cells are >90% blue with X-gal staining).

There was a wide distribution of C17.2 cells (Fig. 1B), suggesting that they had migrated far from their initial sites in the cylinder, compared to the T-10 cells (Fig. 1A), which remained localized to the area of initial seeding in the cylinders. These patterns were observed whether the cells were plated directly on top of the glioma cells (right-sided cylinder [arrows]) or simply in juxtaposition to them (center cylinder [arrows]).

Example 2. Transgene-Expressing NSCs Migrate Throughout and Beyond Invading Tumor Mass *in vivo*

To determine the behavior of clone C17.2 NSCs introduced into brain tumors, experimental animals (syngeneic adult rats) first received an implant of 4×10^4 D74 rat glioma cells in 1 μ l injected into the right frontal lobe. Four days later, 1×10^5 C17.2 NSCs in 1.5 μ l PBS were injected at same coordinates directly into the D74 tumor bed.

25 Animals were then sacrificed at days 2, 6, and 10 post-intratumoral injection and cryostat sections of the brains were processed with Xgal histochemistry for β -galactosidase (β gal) activity to detect donor-derived cells and counterstained with neutral red to detect tumor cells.

Donor C17.2 NSCs were found extensively dispersed throughout the tumor

30 within a few days, spanning an 8mm width of tumor as rapidly as 2 days after injection (Figs.

transduced with the PGK-GFP-IRES-NeoR retroviral vector construct to express green fluorescent protein (GFP) as previously described (ref. Aboody-Guterman et. al, 1997), and maintained in RPMI-1640 (Bio Whittaker) supplemented with 10% FCS and 1% penicillin/streptomycin (Sigma). Cell structure studies were performed in 100mm petri dishes under standard conditions: humidified, 37 ° C, 5% CO₂ incubator. CNS-1 glioma cells were plated to approx. 60-70% confluency around a 5mm cylinder (i.e. free of CNS-1 cells) into which 40,000 C17.2 or TR-10 cells plated overnight. At the same time, 40,000 C17.2 or TR-10 cells were placed into a 5mm cylinder placed directly on top of adhered CNS-1 cells. The next day, cylinders were removed and plates rinsed well with PBS to remove any floating cells, media was replaced, and plates incubated for 5 days. Plates were subsequently stained for β -galactosidase overnight after .5% glutaraldehyde fixation. (Note: both C17.2 and TR-10 cells are >90% blue with X-gal staining).

There was a wide distribution of C17.2 cells (Fig. 1B), suggesting that they had migrated far from their initial sites in the cylinder, compared to the TR-10 cells (Fig. 1A), which remained localized to the area of initial seeding in the cylinders. These patterns were observed whether the cells were plated directly on top of the glioma cells (right-sided cylinder [arrows]) or simply in juxtaposition to them (center cylinder [arrows]).

Example 2. Transgene-Expressing NSCs Migrate Throughout and Beyond Invading Tumor Mass *in vivo*

To determine the behavior of clone C17.2 NSCs introduced into brain tumors, experimental animals (syngeneic adult rats) first received an implant of 4×10^4 D74 rat glioma cells in 1 μ l injected into the right frontal lobe. Four days later, 1×10^5 C17.2 NSCs in 1.5 μ l PBS were injected at same coordinates directly into the D74 tumor bed. Animals were then sacrificed at days 2, 6, and 10 post-intratumoral injection and cryostat sections of the brains were processed with Xgal histochemistry for β -galactosidase (β gal) activity to detect donor-derived cells and counterstained with neutral red to detect tumor cells.

Donor C17.2 NSCs were found extensively dispersed throughout the tumor within a few days, spanning an ~8mm width of tumor as rapidly as 2 days after injection (Figs.

2A, 2B). This is a much more extensive and rapid dispersion compared to previous reports of 3T3 fibroblasts grafted into an experimental brain tumor³³. By day 10, C17.2 cells were seen throughout a majority of the tumor, clearly along the infiltrating tumor edge and slightly beyond it, drawn somewhat by the degenerative environment, seeming to “track” migrating tumor cells (Figs. 2C, 2D). C17.2 cells themselves did not become tumorigenic.

(Fig. 2A) Day 2 shown at 4X; arrowheads demarcate the approximate edges of tumor mass; even at lower power, the tumor can be seen to be intermixed with blue NSCs [arrows]. This is appreciated more dramatically at high power in (Fig. 2B) at 10X where Xgal+, blue-staining NSCs [arrow] are interspersed between tumor cells staining dark red. (Fig. 2C) This view of the tumor mass, 10 days after intra-tumoral injection nicely shows that Xgal+blue, C17.2 NSCs have infiltrated the tumor but largely stop at the edge of the darkly red stained tumor tissue (border indicated by arrowheads) with some migration into surrounding tissue when blue-staining NSC appears to be “following” and invading, “escaping” tumor cell [arrow] (10X). This phenomenon becomes even more dramatic when examining the behavior of C17.2 NSCs in an even more virulent, invasive and aggressive tumor than D74, the experimental CNS-1 astrocytoma in the brain of a nude mouse (Fig. 2D). CNS-1 tumor cells were implanted into an adult nude mouse frontal cortex (day 0). On day 6, 4×10^4 C17.2 cells were implanted directly into the tumor bed. The animal pictured in (Fig. 2D) was sacrificed on day 12 post-tumor implantation, 6 days post-intra-tumoral injection. The cryostat section pictured was processed with Xgal histochemistry for β -galactosidase activity to detect blue C17.2 NSCs and counterstained with neutral red to show dark red tumor cells. There is extensive migration and distribution of blue C17.2 cells throughout the infiltrating experimental tumor bed, up to and along the infiltrating tumor edge [white arrows], and, where many tumor cells are invading normal tissue, into surrounding tissue in virtual juxtaposition to aggressive tumor cells [arrows] (10X).

Example 3. NSCs “Track” Infiltrating Tumor Cells

CNS-1 tumor cells were labelled by retroviral transduction with green fluorescent protein (GFP), prior to implantation, to better distinguish single cells away from the main tumor bed¹⁷. GFP-expressing CNS-1 glioma cells (3×10^4) in 1 μ l PBS injected into right frontal lobe at stereotaxic coordinates 2mm lateral to bregma, on coronal suture, 3mm depth from dura. 4 $\times 10^4$ C17.2 or TR-10 cells in 1 μ l PBS injected at same coordinates directly into tumor bed on day 6. 3-4 C17.2 animals (2 BUdR labelled, 1 BUdR pulsed) and 1-2 TR-10 control animals (1 BUdR labelled). Animals were sacrificed on days 9,12, 16 and 21 post-tumor implantation. Cryostat sectioned, fixed brain tissue was stained either with β -galactosidase (C17.2 cells blue) and neutral red (tumor cells dark red) or double immunofluorescence with Texas red anti- β -galactosidase (C17.2 cells red) and FITC anti-GFP (tumor cells green).

(Figs. 3A, 3B) parallel sections: low power of C17.2 cells distributed throughout tumor and surrounding edge [Fig.3A) Xgal and neutral red, Fig.3B) double immunofluorescent labelling with Texas red and FITC]

(Figs. 3C, 3D) low and high power of single migrating tumor cell in juxtaposition to C17.2 cell (Xgal and neutral red)

(Figs. 3E, 3F) low and high power of single migrating tumor cell in juxtaposition to C17.2 cell (Xgal and neutral red)

(Figs. 3G, 3H) low and high power of single migrating tumor cells in juxtaposition to C17.2 cells (double immunofluorescent labelling with Texas red and FITC).

Example 4. NSCs Implanted at Distant Site Migrate Toward Tumor

To examine the capacity of NSCs to migrate through normal tissue and specifically target tumor cells, donor NSCs were injected into uninvolved sites distant from the main tumor bed in three separate paradigms, into the same hemisphere, into the opposite hemisphere, or into the lateral ventricles.

Same hemisphere: CNS-1 glioma cells (3×10^4) in 1 μ l PBS was injected into the right frontal lobe at stereotaxic coordinates 2mm lateral to bregma, on coronal suture, 3mm depth from dura. 4 $\times 10^4$ C17.2 or TR-10 cells in 1 μ l PBS injected into right frontal parietal lobe at stereotaxic coordinates 3mm lateral and 4mm caudal to bregma, 3mm

depth from dura on day 6. Two animals were sacrificed at days 12 and 21. At all time points, NSCs were found distributed within the main tumor bed as well as in juxtaposition to migrating tumor cells in surrounding tissue (Figs. 4A, 4B).

Opposite hemisphere: 3×10^4 CNS-1 tumor cells in 1 μ l PBS injected into left frontal lobe at stereotaxic coordinates 2mm lateral to bregma, on coronal suture, 3mm depths from dura, 5×10^4 CNS-1 tumor cells in 1 μ l PBS injected into left frontoparietal lobe 3mm lateral and 4mm caudal to bregma, 3mm depth from dura, 8×10^4 C17.2 cells in 2 μ l PBS injected into right frontal lobe 2mm lateral and 2mm caudal to bregma, 3mm depth from dura on day 6. Two animals sacrificed on day 12 and 21. (control - no tumor Coordinates: 2mm R of bregma, 2mm caudal, 3mm deep). NSCs were seen actively migrating across the central commissure towards the tumor on the opposite side of the brain, and then entering the tumor (Figs. 4C, 4D, 4E).

Implantation away from CNS-1 tumor bed (intraventricular):

In this final paradigm 5×10^4 CNS-1 tumor cells in 1 μ l PBS was injected into the right frontal lobe 2mm lateral to bregma, on coronal suture, 3mm depth from dura. 8×10^4 C17.2 cells in 2 μ l PBS injected into left or right ventricle 1mm lateral and 3mm caudal to bregma, 2mm depth from dura on day 6. Two animals sacrificed on days 12 and 21. NSCs again were seen within the main tumor bed, as well as in juxtaposition to migrating tumor cells (Figs. 4F, 4G).

In each case, donor NSCs were found to migrate through normal tissue and “target” the tumor.

11227
 11228
 11229
 11230
 11231
 11232
 11233
 11234
 11235
 11236
 11237
 11238
 11239
 11240
 11241
 11242
 11243
 11244
 11245
 11246
 11247
 11248
 11249
 11250
 11251
 11252
 11253
 11254
 11255
 11256
 11257
 11258
 11259
 11260
 11261
 11262
 11263
 11264
 11265
 11266
 11267
 11268
 11269
 11270
 11271
 11272
 11273
 11274
 11275
 11276
 11277
 11278
 11279
 11280
 11281
 11282
 11283
 11284
 11285
 11286
 11287
 11288
 11289
 11290
 11291
 11292
 11293
 11294
 11295
 11296
 11297
 11298
 11299
 11300
 11301
 11302
 11303
 11304
 11305
 11306
 11307
 11308
 11309
 11310
 11311
 11312
 11313
 11314
 11315
 11316
 11317
 11318
 11319
 11320
 11321
 11322
 11323
 11324
 11325
 11326
 11327
 11328
 11329
 11330
 11331
 11332
 11333
 11334
 11335
 11336
 11337
 11338
 11339
 11340
 11341
 11342
 11343
 11344
 11345
 11346
 11347
 11348
 11349
 11350
 11351
 11352
 11353
 11354
 11355
 11356
 11357
 11358
 11359
 11360
 11361
 11362
 11363
 11364
 11365
 11366
 11367
 11368
 11369
 11370
 11371
 11372
 11373
 11374
 11375
 11376
 11377
 11378
 11379
 11380
 11381
 11382
 11383
 11384
 11385
 11386
 11387
 11388
 11389
 11390
 11391
 11392
 11393
 11394
 11395
 11396
 11397
 11398
 11399
 11400
 11401
 11402
 11403
 11404
 11405
 11406
 11407
 11408
 11409
 11410
 11411
 11412
 11413
 11414
 11415
 11416
 11417
 11418
 11419
 11420
 11421
 11422
 11423
 11424
 11425
 11426
 11427
 11428
 11429
 11430
 11431
 11432
 11433
 11434
 11435
 11436
 11437
 11438
 11439
 11440
 11441
 11442
 11443
 11444
 11445
 11446
 11447
 11448
 11449
 11450
 11451
 11452
 11453
 11454
 11455
 11456
 11457
 11458
 11459
 11460
 11461
 11462
 11463
 11464
 11465
 11466
 11467
 11468
 11469
 11470
 11471
 11472
 11473
 11474
 11475
 11476
 11477
 11478
 11479
 11480
 11481
 11482
 11483
 11484
 11485
 11486
 11487
 11488
 11489
 11490
 11491
 11492
 11493
 11494
 11495
 11496
 11497
 11498
 11499
 11500
 11501
 11502
 11503
 11504
 11505
 11506
 11507
 11508
 11509
 11510
 11511
 11512
 11513
 11514
 11515
 11516
 11517
 11518
 11519
 11520
 11521
 11522
 11523
 11524
 11525
 11526
 11527
 11528
 11529
 11530
 11531
 11532
 11533
 11534
 11535
 11536
 11537
 11538
 11539
 11540
 11541
 11542
 11543
 11544
 11545
 11546
 11547
 11548
 11549
 11550
 11551
 11552
 11553
 11554
 11555
 11556
 11557
 11558
 11559
 11560
 11561
 11562
 11563
 11564
 11565
 11566
 11567
 11568
 11569
 11570
 11571
 11572
 11573
 11574
 11575
 11576
 11577
 11578
 11579
 11580
 11581
 11582
 11583
 11584
 11585
 11586
 11587
 11588
 11589
 11590
 11591
 11592
 11593
 11594
 11595
 11596
 11597
 11598
 11599
 11600
 11601
 11602
 11603
 11604
 11605
 11606
 11607
 11608
 11609
 11610
 11611
 11612
 11613
 11614
 11615
 11616
 11617
 11618
 11619
 11620
 11621
 11622
 11623
 11624
 11625
 11626
 11627
 11628
 11629
 11630
 11631
 11632
 11633
 11634
 11635
 1

REFERENCES

- 17

7. Renfranz PJ, Cunningham MG, McKay RDG : Region-specific differentiation of the hippocampal stem cell line HiB5 upon implantation into the developing mammalian brain. *Cell* 66, 713-729 (1991).
8. Snyder EY, Fisher LJ: Gene therapy for neurologic diseases, *Current Opin. in Pediatrics* 8: 558-568 (1996).
9. Snyder, EY, Taylor, RM, Wolfe, JH: Neural progenitor cell engraftment corrects lysosomal storage throughout the MPS VII mouse brain. *Nature* 374: 367-370 (1995).
10. Lacorazza HD, Flax JD, Snyder EY, Jendoubi, M, Expression of human β -hexosaminidase a-subunit gene (the gene defect of Tay-Sachs disease) in mouse brains upon engraftment of transduced progenitor cells, *Nature Medicine* 4: 424-429 (1996).
11. Lynch WP, Snyder EY, Qualtierre L, Portis JL, Sharpe AH: Neither neurovirulent retroviral envelope protein nor viral particles are sufficient for the induction of acute spongiform neurodegeneration: evidence from engineered neural progenitor-derived chimeric mouse brains, *Journal of Virology* 70: 8896-8907 (1996).
12. Snyder EY, Macklis JD, Multipotent neural progenitor or stem-like cells may be uniquely suited for therapy of some neurodegenerative conditions, *Clinical Neuroscience* 3: 310-316 (1996).
13. Snyder EY, Yoon CH, Flax JD, Macklis JD: Multipotent neural progenitors can differentiate towards the replacement of neurons undergoing targeted apoptotic degeneration in adult mouse neocortex. *Proc Natl Acad Sci* 94:11663-8 (1997).
14. Park KI, Jensen FE, Stieg PE, Snyder EY: Acute CNS injury may direct the migration, proliferation, & differentiation of neural stem cells: Evidence from the effect of hypoxia-ischemia on "reporter" cells, *Soc Neurosci Abstr* 21:2027 (1995).
15. Park KI, Jensen FE, Stieg PE, Himes T, Fisher I, Snyder EY: Transplantation of neurotrophin-3 (NT-3) expressing neural stem-like cells into hypoxic-ischemic (HI) brain injury, *Soc Neurosci Abstr* 23: 346 (1997).

16. Park KI, Liu S, Flax JD, Nissim S, Stieg PE, Snyder EY, Transplantation of neural progenitor & stem-like cells: developmental insights may suggest new therapies for spinal cord and other CNS dysfunction, *Journal of Neurotrauma* (in press).
17. Aboody-Guterman KS, Pechan PA, Rainov NG, Sena-Esteves M, Snyder EY, Wild P, Schraner E, Tobler K, Breakefield XO and Fraefel C: Green fluorescent protein as a reporter for retrovirus and helper virus-free HSV-1 amplicon vector-mediated gene transfer into neural cells in culture and *in vivo*. *NeuroReport* Vol 8:3801-8 (1997).
18. Kruse CA, Molleston M, Parks EP, Schiltz PM, Kleinschmidt-DeMasters BK, and Hickey WF: A rat glioma model, CNS-1, with invasive characteristics similar to those of human gliomas: A comparison to 9L gliosarcoma. *J of Neuro-Oncology* 22:191-200 (1994).
19. Short et al
20. Taniya T, Wei, MX, Chase M, Ono Y, Lee F, Breakefield XO, and Chiocca EA: Transgene inheritance and retroviral infection contribute to the efficiency of gene expression in solid tumors inoculated with retroviral vector producer cells. *Gene therapy* 2:531-538, (1995).
21. Deonarain MP, Spooner RA, and Epenetos AA: Genetic delivery of enzymes for cancer therapy. *Gene Therapy* 2:235-244 (1995).
22. Kramm CM, Sena-Esteves M, Barnett F, Rainov N, Schuback D, Yu J, Pechan P, Paulus W, Chiocca EA, and Breakefield XO: Gene therapy for brain tumors. *Brain Pathology*, 5:345-381 (1995).
23. Philpott GW, Shearer WT, Bower RW and Parker CW: Selective cytotoxicity of hepten-substituted cells with an antibody-enzyme conjugate. *J Immunol.* 111:921-929 (1973).
24. Huber BE, Austin EA, Richards CA, Davis ST and Good S: Metabolism of 5-fluorocytosine to 5-fluorouracil in human colorectal tumor cells transduced with the cytosine deaminase gene: Significant antitumor effects when only a small percentage of tumor cells express cytosine deaminase. *Proc. Natl. Acad Sci.* 91:8302-8306 (1994).

25. Mullen C, Kilstrup M, and Blaese RM: Transfer of the bacterial gene for cytosine deaminase to mammalian cells confers lethal sensitivity to 5-fluorocytosine: a negative selection system. *Proc. Natl. Sci. USA* 89:33-37 (1992.)
26. Kun Le, Gajjar A, Muhlbaue M et al: Clinical protocol: stereotactic injection of herpes simplex thymidine kinase vector producer cells (PA317-GITk1SvNa.7) and intravenous gangciclovir for the treatment of progressive or recurrent primary supratentorial pediatric malignant brain tumors, *Hum Gene Ther* 6: 1231-1255 (1995).
27. Walther W, Stein U, Pfeil D: Gene transfer of human TNF α into glioblastoma cells permits modulation of *mdr1* expression and potentiation of chemosensitivity. *Int J Cancer* 61:832-839 (1995).
28. Manome y, Wen PY, Dong Y et al: Viral vector transduction of the human deoxycytidine kinase cDNA sensitizes glioma cells to the cytotoxic effects of cytosin arabinoside *in vitro* and *in vivo*. *Nat Med* 2:567-573 (1996).
29. Takamiya Y, Short MP, Moolten F, Fleet C, Mineta T, Breakefield XO, Martuza RL: An experimental model of retroviral gene therapy for malignant brain tumors, *J Neurosurg* 79: 104-110 (1993).
30. Barba D, Hardin J, Ray J, Gage FH : Thymidine kinase-mediated killing of rat brain tumors, *J Neurosurg* 79: 729-735 (1993).
31. Vrionis FD, Wu JK, QI P, Cano WG, Cherington V Tumor cells expressing the herpes simplex virus-thymidine kinase gene in the treatment of Walker 256 meningeal neoplasia in rats. *J Neurosurg* 84:250-257 (1996).
32. Lyons RM, Forry-Schaudies S, Otto E, et al: An improved retroviral vector encoding the herpes simplex thymidine kinase gene increases antitumor efficacy *in vivo*. *Cancer Gene Ther* 2:273-280 (1995).
33. Rainov NG, Kramm CM, Aboody-Guterman K, Chase M, Ueki K, Louis D, Harsh G, Chiocca EA, and Breakefield XO: Retrovirus-mediated gene therapy of experimental brain

neoplasms using the herpes simplex virus-thymidine kinase/ganciclovir paradigm. *Cancer Gene Therapy* Vol 3:2:99-106 (1996).

34. Somia N, Zoppe M, Verma IM Generation of targeted retroviral vectors by using single-chain variable fragment: An approach to *in vivo* gene delivery. *Proc Natl Acad Sci USA* 92:7570-7574 (1995).
35. Chen C, Chang Y, Ryan P et al (1995): Effect of herpes simplex virus thymidine kinase expression level on ganciclovir-mediated cytotoxicity and the "bystander effect". *Hum Gene Ther* 6: 1467-1476.
36. Weinstein DE et al: C17 inhibits the proliferation of astrocytes & astrocytoma cells by a contact-mediated mechanism, *Glia* 3: 130-139 (1990).